Flemingins G–O, Cytotoxic and Antioxidant Constituents of the Leaves of *Flemingia grahamiana*Ivan Gumula,^{†,‡} John Patrick Alao,[‡] Isaiah Omolo Ndiege,[§] Per Sunnerhagen,[‡] Abiy Yenesew,^{*,†} and Máté Erdélyi^{*,‡,⊥}[†]Department of Chemistry, University of Nairobi, POB 30197-00100, Nairobi, Kenya[‡]Department of Chemistry and Molecular Biology, University of Gothenburg, SE-41296 Gothenburg, Sweden[§]Office of the Vice Chancellor, Kyambogo University, POB 1, Kyambogo-Kampala, Uganda[⊥]Swedish NMR Centre, University of Gothenburg, SE-40530 Gothenburg, Sweden

S Supporting Information

ABSTRACT: The known flemingins A–C (1–3) and nine new chalcones, named flemingins G–O (4–12), along with deoxyhomoflemingin (13) and emodin (14) were isolated from a leaf extract of *Flemingia grahamiana*. The isolated chalcones were found to have a geranyl substituent modified into a chromene ring possessing a residual chain, as shown by spectroscopic methods. The leaf extract showed an IC₅₀ value of 5.9 μg/mL in a DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging assay. The chalcones flemingins A, B, C, G, and H were active in the DPPH radical scavenging assay (ED₅₀ 4.4–8.9 μM), while flemingins A and C showed cytotoxicity against MCF-7 human breast cancer cells (IC₅₀ 8.9 and 7.6 μM, respectively).

Flemingia grahamiana Wight & Arn. (Leguminosae) is a shrub or semishrub reaching a height of up to 1.8 m. It is native to tropical Africa, occurring in open and wooded savannas, sometimes near water in riverine vegetation, on hillsides, in termite mounds, and along roadsides.^{1,2} Its fruits and inflorescence are the sources of *warrus* (*warus*), a common cosmetic dye in East Africa, Saudi Arabia, and India.^{2,3} This dye has also been obtained from other *Flemingia* species.³ The roots of *F. grahamiana* are used in Eastern and Southern Africa for the treatment of diarrhea and dysentery, whereas in India the leaves of the plant are used as a purgative and for the treatment of skin diseases.² Cardillo et al.^{3,4} have analyzed its constituents, from a sample collected in Eritrea, leading to the identification of flemingins A–F, homoflemingin, and deoxyhomoflemingin. Despite its wide use, the biological activities of *F. grahamiana* extracts and constituents have not yet been studied. As part of an ongoing search for novel bioactive agents, the isolation, identification, and the cytotoxic and antioxidant activities of chalcones from the leaves of *F. grahamiana* are reported herein.

RESULTS AND DISCUSSION

Column chromatographic fractionation of a CH₂Cl₂–CH₃OH (1:1) extract of the leaves of *F. grahamiana* with subsequent purification by gel filtration over Sephadex LH-20 and by preparative HPLC led to the isolation of 14 secondary metabolites, including the known flemingins A (1), B (2), and C (3)^{3,4} and nine new chalcones (4–12), together with deoxyhomoflemingin^{4,5} (13) and emodin⁶ (14). Compounds

1–14 were identified using spectroscopic data interpretation. The identities of the known compounds, 1–3, 13, and 14, were confirmed by comparison of their spectroscopic data to those previously published (Tables S1 and S2 and spectra S1–S19 and S99–S07, Supporting Information).^{3–6}

Compound 4, a yellow solid, was obtained as a diastereomeric mixture. Analysis of compound 4 by HRESIMS gave a pseudomolecular ion peak at *m/z* 439.1771 [*M* + *H*]⁺, corresponding to the molecular formula C₂₅H₂₇O₇ ([*M* + *H*]⁺, calcd 439.1757). Its NMR spectra (Tables 1 and 2) resembled those of 1–3 (Tables S1 and S2) and were thus indicative of a chalcone substituted with a geranyl group modified into a chromene ring possessing a side chain, which is a common feature of the secondary metabolites of the genus *Flemingia*. Its ¹H NMR spectrum exhibited a pair of doublets at δ_H 8.06 (1H, H-β) and 7.72 (1H, H-α) with the vicinal coupling constant ³*J*_{HH} = 15.2 Hz indicative of their *trans* relationship. The corresponding carbons resonating at δ_C 141.7 (C-β) and 121.3 (C-α) were identified from the HSQC spectrum (Table 2). Both H-α (δ_H 7.72) and H-β (δ_H 8.06) showed HMBC cross-peaks to the carbonyl carbon at δ_C 194.0, suggesting that they are part of an α,β-unsaturated ketone moiety of a chalcone. The ¹H NMR signals of the A-ring protons of 4 gave a two-proton multiplet at δ_H 6.74 arising from the strongly coupled H-3 and

Received: May 21, 2014

Published: September 16, 2014

Table 1. ^1H NMR Spectroscopic Data for Flemingins G–J (4–7) (δ_{H} , multiplicity (J in Hz))^a

position	4	5	6	7
3	6.74, m	6.37, d (8.2)	6.93, d (8.0)	6.95, d (8.0)
4	6.74, m	7.02, dd (8.2, 8.2)	7.27, dd (8.0, 7.2)	7.28, dd (8.0, 7.2)
5		6.37, d (8.2)	6.87, dd (8.0, 7.2)	6.88, dd (8.0, 7.2)
6	7.03, dd (1.8, 1.8)		7.83, d (8.0)	7.84, d (8.0)
β	8.06, d (15.2)	8.37, d (15.6)	8.06, d (15.2)	8.07, d (15.2)
α	7.72, d (15.2)	8.14, d (15.6)	7.79, d (15.2)	7.80, d (15.2)
6'	7.33, s	7.28, s	7.43, s	7.49, s
3''	5.63/5.63, d (10.0/10.4)	5.62/5.62, d (10.0/10.0)	5.68, d (9.6)	5.70, d (9.6)
4''	6.76/6.76, d (10.4/10.0)	6.76/6.76, d (10.0/10.0)	6.63, d (9.6)	6.65, d (9.6)
5''	1.46, s	1.46, s	1.38, s	1.39, s
6''a	1.63/1.78, m	1.63/1.77, m	1.63, m	1.53, m
6''b	1.78/1.89, m	1.78/1.89, m	1.68, m	1.76, m
7''a	1.66/1.71, m	1.65/1.67, m	1.22, m	1.24, m
7''b	1.71/1.78, m	1.72/1.74, m	1.50, m	1.53, m
8''	3.99/4.01, m	3.99/4.01, m	3.85, m	3.85, m
10''a	4.80, dd (5.4, 1.0)	4.80, ddd (5.3, 1.8, 1.8)	4.70, m	4.72, m
10''b	4.91, dd (5.4, 1.4)	4.91, m	4.83, m	4.85, m
11''	4.68, dd (1.4, 1.0)	1.69, m	1.59, m	1.60, m
OH-2	n.o. ^b	n.o. ^b	10.34, s	10.35, s
OH-2'	n.o. ^b	n.o. ^b	13.54, s	13.55, s
OH-5'	n.o. ^b	n.o. ^b	8.70, s	8.72, s
OH-8''	n.o. ^b	n.o. ^b	10.19, s	10.2, s

^a ^1H NMR spectra for compounds 4 and 5 were acquired in CD_3OD , while those for compounds 6 and 7 were obtained in $\text{DMSO}-d_6$ at 25 °C. ^bn.o., not observed.

H-4, and a doublet of doublets ($^4J_{\text{HH}}$; $^5J_{\text{HH}}$ = 1.8, 1.8 Hz) integrating for one proton at δ_{H} 7.03 (H-6), the spin system of which was confirmed using the COSY spectrum. The assignment was supported by the HMBC cross-peaks of H-6 to C- β ($^3J_{\text{CH}}$ to δ_{C} 141.7), C-2 ($^3J_{\text{CH}}$ to δ_{C} 152.3), and C-4 ($^3J_{\text{CH}}$ to δ_{C} 120.5). The HMBC cross-peaks H- β to C-2 ($^3J_{\text{CH}}$ to δ_{C} 152.3) and C-6 ($^3J_{\text{CH}}$ to δ_{C} 115.6) provided confirmation of the nature of the A-ring. A NOE cross-peak between H- α (δ_{H} 7.72) and a singlet at δ_{H} 7.33 (Figure 1) allowed the assignment of this signal to H-6', which is the only proton in the B-ring.

The substitution pattern of the B-ring of compound 4, with a chromene at C-3' and C-4' formed by cyclization of a geranyl group, and two hydroxy groups at C-2' and C-5', was found to be similar to those of compounds 1–3,³ the only exception being the nature of the C-2'' substituent. The assignment of the singlet at δ_{H} 7.33, 1H, to H-6' was confirmed by its HMBC correlations to three quaternary aromatic carbons, C-2' ($^3J_{\text{CH}}$ to δ_{C} 155.7), C-4' ($^3J_{\text{CH}}$ to δ_{C} 150.1), and C-5' ($^2J_{\text{CH}}$ to δ_{C} 138.9), and the carbonyl carbon, C=O ($^3J_{\text{CH}}$ to δ_{C} 194.0). The low chemical shift of C-5' could be attributed to the shielding effect of oxygenation at C-2' and C-4'. Duplicate signals of two *ortho* olefinic protons at δ_{H} 5.63/5.63 (1H, d, 3J = 10.1/10.4 Hz) and 6.76/6.76 (1H, d, 3J = 10.1/10.4 Hz) were assigned to H-3'' and H-4'', respectively, of the chromene ring, for which the placement at C-3'/C-4' was defined by the HMBC cross-peaks

of H-3'' to C-3' ($^3J_{\text{CH}}$ to δ_{C} 110.7) and of H-4'' to C-2' ($^2J_{\text{CH}}$ to δ_{C} 155.7) and C-4' ($^3J_{\text{CH}}$ to δ_{C} 150.1). The deduction that the side chain of compound 4 has a terminal double bond at C-9'' (δ_{C} 148.6) and a hydroxy group at C-8'' (δ_{C} 76.5/76.6) was based on the signal at δ_{H} 3.99–4.01 (1H, m) attributed to H-8'', the multiplets at δ_{H} 4.80 (1H) and 4.91 (1H) due to the H-10''a (1H, dd) and 10''b (1H, dd) methylene protons of the terminal double bond, and the allylic methyl singlet at δ_{H} 1.68 (3H, H-11''). The latter proton signal showed HMBC correlations to C-8'' ($^3J_{\text{CH}}$ to δ_{C} 76.6/76.5), C-10'' ($^3J_{\text{CH}}$ to δ_{C} 111.4/111.7), and C-9'' ($^2J_{\text{CH}}$ to δ_{C} 148.6). Moreover, H-6''a/b (δ_{H} 1.63/1.78, 1H and 1.78/1.89, 1H) exhibited $^3J_{\text{CH}}$ correlations to C-3'' (δ_{C} 128.2) and C-5'' (δ_{C} 27.4/27.6), indicating that C-2'' is the position of attachment of the substituent on the chromene ring. Assignment of the 10''a and 10''b protons (δ_{H} 4.80, 1H and 4.91, 1H) was made based on the observation of a NOE correlation between H-10''a and CH_3 -11'' (δ_{H} 1.68, 3H), as well as one between H-10''b (δ_{H} 4.91, 1H) and H-8'' (δ_{H} 3.99/4.01, 1H) (Figure 1). The signal duplications, observed in the ^1H and ^{13}C NMR spectra, for positions 3'', 4'', 6'', and 8'' indicated that compound 4 was isolated as a diastereomeric mixture, with its stereocenters being at C-2'' and C-8''. This new compound was therefore characterized as 2,5,2',5'-tetrahydroxy-2''-(3'''-hydroxy-4'''-methylpent-4'''-enyl)-2-methylpyrano-[5'',6'':3',4']-chalcone and was assigned the trivial name *flemingin G*.

Compound 5 was identified as a diastereomeric mixture of a chalcone. Its NMR spectra (Tables 1 and 2) showed close similarities to those of compound 4, except for the C-2 and C-6 oxygenation of its A-ring, providing an AX_2 spin system with δ_{H} 7.02 (1H, dd, 3J = 8.4, 8.2 Hz, H-4) and 6.37 (2H, d, 3J = 8.2 Hz, H-3/5). Its HRESIMS peak at m/z 439.1809 $[\text{M} + \text{H}]^+$ was consistent with the molecular formula $\text{C}_{25}\text{H}_{27}\text{O}_7$ ($[\text{M} + \text{H}]^+$, calcd 439.1757). On the basis of the above data this new compound, *flemingin H*, was characterized as 2,6,2',5'-tetrahydroxy-2''-(3'''-hydroxy-4'''-methylpent-4'''-enyl)-2-methylpyrano-[5'',6'':3',4']-chalcone.

Compounds 6 and 7 eluted as distinct fractions by preparative HPLC and were identified by NMR as being diastereomers. Their NMR spectra indicated a chalcone core with identical B-ring substitution and the same C-2'' substituent as those of compounds 4 and 5, and a C-2 monohydroxylated A-ring analogous to *flemingin A* 1.³ Hence, the A-ring of compound 6 gave rise to a doublet of doublets at δ_{H} 6.87 (1H, 3J = 8.0, 7.2 Hz, H-5), a doublet at δ_{H} 6.93 (1H, 3J = 8.0 Hz, H-3), a doublet of doublets at δ_{H} 7.27 (1H, 3J = 8.0, 7.2 Hz, H-4), and a doublet at δ_{H} 7.83 (1H, 3J = 8.0, Hz), attributable to H-6.

The C-2 oxygenation of this ring was supported by HMBC correlations of H-4 (δ_{H} 7.27, 1H) to the quaternary carbon C-2 ($^3J_{\text{CH}}$ to δ_{C} 157.4) and to the aromatic methine carbon C-6 ($^3J_{\text{CH}}$ to δ_{C} 129.3). Moreover, H-3 (δ_{H} 6.93, 1H), H-6 (δ_{H} 7.83, 1H), and H- α gave $^3J_{\text{CH}}$ correlations to the quaternary carbon C-1 (δ_{C} 121.3), and H- β to C-2 (δ_{C} 157.4) and C-6 (δ_{C} 129.3).

Compound 7 showed virtually identical NMR spectra to 6, with very minor differences in chemical shifts. The largest difference in chemical shift between the ^1H NMR signals of compounds 6 and 7 was observed for the H-6' of their B-ring, resonating at δ_{H} 7.43 and 7.49, respectively. The assignment of these signals was confirmed by the $^3J_{\text{CH}}$ HMBC correlation of H-6' to the C=O moiety (δ_{C} 192.0), for both compounds 6 and 7. Furthermore, H-6''a and H-6''b resonated at δ_{H} 1.63 and 1.68, respectively, for compound 6, but at δ_{H} 1.53 and 1.76 for compound 7. The very similar chemical shifts and the identical

Table 2. ^{13}C NMR Spectroscopic Data for Compounds 4–7^a

position	4	5	6	7
1	123.5, C	111.5, C	121.3, C	121.3, C
2	152.3, C	160.6, C	157.4, C	157.4, C
3	117.7, CH	107.8, CH	116.3, CH	116.3, CH
4	120.5, CH	132.9, CH	132.2, CH	132.2, CH
5	151.3, C	107.8, CH	119.5, CH	119.5, CH
6	115.6, CH	160.6, C	129.3, CH	129.3, CH
β	141.7, CH	138.0, CH	139.6, CH	139.6, CH
α	121.3, CH	122.4, CH	119.8, CH	119.8, CH
C=O	194.0, C	195.2, C	192.0, C	192.0, C
1'	113.8, C	114.1, C	112.1, C	112.1, C
2'	155.7, C	155.6, C	153.8, C	153.8, C
3'	110.7, C	110.7, C	108.9, C	109.0, C
4'	150.1, C	149.7, C	149.0, C	148.9, C
5'	138.9, C	138.8, C	137.9, C	137.9, C
6'	116.0, CH	116.0, CH	115.4, CH	115.4, CH
2''	81.7/81.8, C	81.5/81.7, C	80.2, C	80.1, C
3''	128.3 CH	128.2, CH	127.8, CH	127.9, CH
4''	118.0, CH	117.8, CH	115.9, CH	115.9, CH
5''	27.4/27.6, CH ₃	27.4/27.5, CH ₃	26.7, CH ₃	26.6, CH ₃
6''	38.5/38.5, CH ₂	38.5/38.5, CH ₂	36.9, CH ₂	36.9, CH ₂
7''	30.2/30.3, CH ₂	30.2/30.4, CH ₂	29.0, CH ₂	29.0, CH ₂
8''	76.5/76.6, CH	76.5/76.6, CH	73.7, CH	73.8, CH
9''	148.6, C	148.6, C	147.9, C	147.8, C
10''	111.4/111.7, CH ₂	111.7, CH ₂	110.1, CH ₂	110.2, CH ₂
11''	17.5/17.7, CH ₃	17.5/17.7, CH ₃	17.6, CH ₃	17.5, CH ₃

^a ^{13}C NMR spectra for compounds 4 and 5 were acquired in CD_3OD , and for compounds 6 and 7 in $\text{DMSO}-d_6$ (25 °C).

Figure 1. Key NOE correlations (blue arrows) observed for compound 4 (mixing time 700 ms, CD_3CN , 25 °C, 800 MHz, Spectrum S23, Supporting Information).

coupling pattern of the NMR signals of 6 and 7 (Tables 1 and 2) suggested that these compounds are diastereomers. The HRESIMS pseudomolecular ion at m/z 423.1821 and 423.1843 $[\text{M} + \text{H}]^+$, observed for 6 and 7, respectively, were consistent with the same molecular formula $\text{C}_{25}\text{H}_{27}\text{O}_6$ ($[\text{M} + \text{H}]^+$, calcd 423.1808). The CD spectra of these compounds, acquired in methanol, did not show significant Cotton effects, indicating both substances to be racemic. On the basis of these data, compounds 6 and 7 were characterized as 2,2',5'-trihydroxy-2''-(3'''-hydroxy-4'''-methylpent-4'''-enyl)-2''-methylpyrano-[5'',6'':3',4']-chalcone diastereomers. These new compounds, for which the diastereomers were not distinguished, were given the trivial names flemingins I and J, respectively.

Compound 8 was obtained as a yellow solid and was assigned a molecular formula of $\text{C}_{25}\text{H}_{26}\text{O}_7$ based on analysis of its HRESIMS ($\text{C}_{25}\text{H}_{26}\text{O}_7$, $[\text{M} + \text{H}]^+$ m/z 439.1733, calcd 439.1757). This substance was identified as being a chalcone derivative with NMR spectroscopic features very similar to those of compound 4, but with an aliphatic chain attached to C-2'', isomeric to that of 4, and thus carrying a hydroxy group at C-9'' and a double bond between C-7'' and C-8''. This double bond was identified by the COSY correlations of the diastereotopic protons δ_{H} 2.42 (1H, m, H-6''a) and 2.49

(1H, m, H-6''b) with the olefinic protons resonating at δ_{H} 5.68 (2H, m, H-7'' and H-8'', Table 4). The assignment was supported by the HMBC correlation of H-6'' (δ_{H} 2.42 and 2.49) and the olefinic carbon at δ_{C} 121.8 (C-7'') and of the two methyl groups CH_3 -10'' and CH_3 -11'' (δ_{H} 1.16, 3H and δ_{H} 1.17, 3H) to the sp^2 methine carbon at δ_{C} 143.4 (C-8'', $^3J_{\text{CH}}$) as well as to an sp^3 oxygenated quaternary carbon at δ_{C} 71.2 (C-9'', $^2J_{\text{CH}}$). As the $^3J_{\text{HH}}$ was not readable due to the identical chemical shift of H-7'' and H-8'', the proposed *E*-configuration of the C-7''–C-8'' double bond was based on the chemical shift of its carbons (predicted values for *E*: $\delta_{\text{C}-8''}$ 142 and $\delta_{\text{C}-7''}$ 124, and for *Z*: $\delta_{\text{C}-8''}$ 137 and $\delta_{\text{C}-7''}$ 120; observed $\delta_{\text{C}-8''}$ 143.4 and $\delta_{\text{C}-7''}$ 121.8).⁷ While this assignment is less reliable than that based on the magnitude of the $^3J_{\text{HH}}$ of the olefinic protons, it is in good agreement with conclusions made by biogenetic considerations.⁸ This compound, flemingin K, was therefore characterized as 2,5,2',5'-tetrahydroxy-2''-((*E*)-4'''-hydroxy-4'''-methylpent-2'''-enyl)-2''-methylpyrano-[5'',6'':3',4']-chalcone (8).

Compound 9 was isolated as a yellow solid. The HRESIMS at m/z 423.1832 $[\text{M} + \text{H}]^+$ was compatible with the molecular formula, $\text{C}_{25}\text{H}_{27}\text{O}_6$ ($[\text{M} + \text{H}]^+$, calcd 423.1808). On the basis of spectroscopic similarities to 4–8, compound 9 was also identified as a chalcone. The spectroscopic features of the B- and C-rings, as well as for the C-2'' substituent, were virtually identical to those of compounds 8 and 9 (Tables 3 and 4). However, compound 9 was found to possess a 2-hydroxyated A-ring, as revealed by its coupling pattern, exhibiting a multiplet at δ_{H} 6.88 (1H, H-3), a doublet of doublets of doublets at δ_{H} 7.25 (1H, $J = 7.8, 7.8, 2.0$ Hz, H-4), a multiplet at δ_{H} 6.90 (1H, H-5), and a doublet of doublets centered at δ_{H} 7.63 (1H, $J = 8.0, 2.0$ Hz, H-6). Although the scalar coupling of H-7'' and H-8'' was not measurable due to their similar shifts in CD_3OD , its

Table 3. ^{13}C NMR Spectroscopic Data for Compounds 8–12^a

position	8	9	10	11	12
1	123.5, C	121.3, C	121.3, C	123.2, C	121.4, C
2	152.3, C	157.0, C	157.4, C	159.0, C	157.3, C
3	118.0, CH	116.1, CH	115.9, CH	117.1, CH	116.3, CH
4	120.5, CH	132.2, CH	132.2, CH	132.9, CH	132.1, CH
5	151.4, C	119.5, CH	119.5, CH	120.9, CH	119.5, CH
6	115.7, CH	129.3, CH	129.3, CH	131.1, CH	129.3, CH
β	141.7, CH	139.6, CH	139.6, CH	141.8, CH	139.2, CH
α	121.3, CH	119.4, CH	119.8, CH	121.5, CH	120.1, CH
C=O	194.0, C	191.2, C	191.9, C	194.2, C	192.0, C
1'	113.9, C	112.1, C	115.3, C	113.9, C	112.7, C
2'	155.7, C	153.8, C	153.8, C	155.8, C	158.6, C
3'	111.0, C	110.0, C	109.0, C	110.8, C	111.3, C
4'	150.2, C	148.9, C	149.0, C	150.1, C	149.8, C
5'	138.9, C	137.9, C	138.0, C	139.0, C	138.8, C
6'	115.9, CH	115.5, CH	112.0, CH	116.0, CH	111.9, CH
1''					31.0, CH
2''	81.5, C	79.9, C	80.5, C	81.8, C	121.6, CH
3''	128.1	127.4, CH	128.0, CH	128.7, CH	133.6, C
4''	117.8, CH	116.3, CH	116.3, CH	117.7, CH	28.9, CH ₂
5''	27.2, CH ₃	26.3, CH ₃	29.0, CH	27.2, CH ₃	20.3, CH ₂
6''	45.3, CH ₂	43.4, CH ₂	38.6, CH ₂	39.8, CH ₂	38.7, CH
7''	121.8, CH	119.8, CH	26.7, CH ₂	26.7, CH ₂	78.1, C
8''	143.4, CH	143.3, CH	77.6, CH	79.7, CH	24.9, CH ₃
9''	71.2, C	68.8, C	71.6, C	73.9, C	25.2, CH ₃
10''	29.6, CH ₃	29.83, CH ₃	24.4, CH ₃	24.9, CH ₃	23.5, CH ₃
11''	29.7, CH ₃	29.83, CH ₃	25.4, CH ₃	25.8, CH ₃	

^a ^{13}C NMR spectra for compounds 8, 9, and 11 were in CD₃OD, and for compounds 10 and 12 in DMSO-*d*₆ (25 °C).

Table 4. ^1H NMR Spectroscopic Data for Flemingins K–M (8–10) (δ_{H} , multiplicity (J in Hz))^a

position	8	9	10
3	6.74, m	6.88, m	6.94, d (8.3)
4	6.74, m	7.25, ddd (7.8, 7.8, 2.0)	7.28, ddd (8.3, 7.3, 1.6)
5		6.90, dd (7.8, 8.0)	6.89, dd (8.0, 7.3)
6	7.03, dd (1.7, 1.7)	7.63, dd (8.0, 2.0)	7.84, dd (8.0, 1.6)
β	8.05, d (15.2)	8.11, d (15.6)	8.08, d (15.6)
α	7.72, d (15.2)	7.82, d (15.6)	7.81, d (15.6)
6'	7.32, s	7.34, s	7.49, s
3''	5.64, d (9.6)	5.64, d (10.2)	5.69, d (10.1)
4''	6.76, d (9.6)	6.76, d (10.2)	6.65, d (10.1)
5''	1.49, s	1.49, s	1.39, s
6''a,b	2.42, m	2.44, m	1.68, m
	2.49, m	2.47, m	1.95, m
7''a,b	5.68, m	5.68, m	1.23, m
			1.68, m
8''	5.68, m	5.68, m	3.04, dd (10.1, 6.1)
10''	1.16, s	1.16, s	0.96, s
11''	1.17, s	1.17, s	1.03, s

^a ^1H NMR spectra for compounds 8–10 were acquired in CD_3OD , and for compound 11 in $\text{DMSO}-d_6$ (25 °C).

magnitude determined in $\text{DMSO}-d_6$ ($^3J_{\text{HH}} = 10.2$ Hz) was indicative of a *trans* configuration. This new compound, flemingin L, was therefore identified as 2,2',5'-trihydroxy-2''-((*E*)-4'''-hydroxy-4'''-methylpent-2'''-enyl)-2''-methylpyrano-[5'',6'':3',4']-chalcone.

Compounds 10 and 11 were obtained by HPLC separation as yellow solids. These compounds were identified by NMR as diastereomers of a chalcone with an identical 2''-substituent to compounds 10 and 11 and possess a C-2-oxygenated A-ring similar to 1,³ 6, and 7. The assignment of the ABCD spin system of the A-ring was based on COSY and TOCSY spectra and was supported by the NOE interaction of H- β (δ_{H} 8.08/8.11) and H-6 (δ_{H} 7.84/7.63). Apart from some small differences in chemical shift for H-6'' and H-7'' (Tables 4 and 5), compounds 10 and 11 showed almost superimposable ^1H NMR spectra (Figure 2). Their HRESIMS gave an $[\text{M} + \text{H}]^+$ peak at m/z 441.1779 and 441.1912 for 10 and 11, respectively, corresponding to the molecular formula $\text{C}_{25}\text{H}_{29}\text{O}_7$ (calcd 441.1913).

The $^3J_{\text{CH}}$ HMBC correlation of H-6' (δ_{H} 1.68/1.73 and 1.95/2.12) with C=O (δ_{C} 191.9/194.2), C-2' (δ_{C} 153.8/155.8), and C-4' (δ_{C} 149.0/150.1) and its $^2J_{\text{CH}}$ correlation to C-5' (δ_{C} 138.0/139.0) indicated that these compounds are diastereomers and not regioisomers. Due to the use of $\text{DMSO}-d_6$ as solvent, the chelated hydroxy group OH-2' could be identified (δ_{H} 13.6). From the above data, these two new compounds, named flemingins M and N, were identified as diastereomers of 2,2',5'-trihydroxy-2''-(3'',4''-dihydroxy-4'''-methylpentyl)-2''-methylpyrano-[5'',6'':3',4']-chalcone. Their CD spectra showed insignificant Cotton effects, thus indicating that both are racemic at C-2''.

Compound 12 was also obtained as a yellow solid and was identified as a chalcone derivative possessing a monooxygenated A-ring, which was revealed by the similar ^1H NMR chemical shifts and COSY spectrum to those of compounds 1, 6, 7, 10, and 11. The nature of the A-ring was confirmed by the presence of only one carbon, C-2, in this ring with a chemical shift above 150 ppm in the ^{13}C NMR spectrum (Table 3) and

Table 5. ^1H NMR Spectroscopic Data for Flemingins N and O (11 and 12) (δ_{H} , multiplicity (J in Hz))^a

position	11	12
3	6.89, m	6.94, dd (8.2, 1.2)
4	7.25, ddd (8.7, 7.3, 1.6)	7.28, ddd (8.2, 7.5, 1.6)
5	6.90, m	6.88, ddd (7.9, 7.5, 1.6)
6	7.63, dd (8.0, 1.6)	7.84, dd (7.9, 1.6)
β	8.11, d (15.5)	8.06, d (15.6)
α	7.82, d (15.5)	7.81, d (15.6)
6'	7.35, s	7.40, s
1''		3.56, m
2''		6.26, m
3''	5.67, d (10.2)	
4''a	6.76, d (10.2)	1.91, m
4''b		1.98, m
5''a	1.48, s	1.27, m
5''b		1.92, m
6''a	1.73, m	1.83, m
6''b	2.12, m	
7''a	1.45, m	
7''b	1.81, m	
8''	3.25, dd (10.8, 2.0)	1.25, s
9''		1.41, s
10''	1.12, 3H, s	1.63, s
11''	1.15, s	

^a ^1H NMR spectrum for compound 11 was obtained in CD_3OD , and for compound 12 in $\text{DMSO}-d_6$ (25 °C).

also by the NOE correlation of H- β (δ_{H} 8.06, 1H) and H-6 (δ_{H} 7.81, 1H) (Figure 3).

Unlike chalcones 1–11, where the C-ring has a residual side chain at C-2'', the side chain of compound 12 was found to be fused with a methylcyclohexene ring. Thus, the ^{13}C NMR spectrum of 12 showed seven aliphatic carbons, two of them tertiary CHs at δ_{C} 31.0 (C-1'') and 38.7 (C-6''), two methylenes at δ_{C} 20.3 (C-5'') and 28.9 (C-4''), and three methyl carbons at δ_{C} 23.5 (C-10''), 24.9 (C-8''), and 25.2 (C-9''). This could be attributed either to a highly saturated geranyl chain or a cyclohexene ring system, with the latter deduced from the COSY correlation of the benzylic/allylic proton δ_{H} 3.56 (1H, H-1'') to the olefinic proton at δ_{H} 6.26 (1H, H-2'') and to the aliphatic proton at δ_{H} 1.83 (1H, H-6''). The presence of a cyclohexene ring in 12 was supported by the HMBC correlations of H-10'' (δ_{H} 1.63, 3H) to the sp^2 carbons at δ_{C} 121.6 (C-2'') and 133.6 (C-3'') and to the sp^3 methylene carbon at δ_{C} 28.9 (C-4'') and the HMBC correlations of CH_3 -8''/9'' (δ_{H} 1.25, 3H and δ_{H} 1.41, 3H) and C-6'' (δ_{C} 38.7). The *cis* configuration of the dihydropyran-cyclohexene ring junction was established from the NOE interaction between H-1'' (δ_{H} 3.56, 1H) and H-6'' (δ_{H} 1.83, 1H). On the basis of the above data, 12 was identified as (*E*)-1-((6a,10a)-1,4-dihydroxy-6,6,9-trimethyl-6a,7,8,10a-tetrahydro-6H-benzo[*c*]chromen-2-yl)-3-(2-hydroxyphenyl)prop-2-en-1-one and was given the trivial name flemingin O.

Motivated by the use of *F. grahamiana* in traditional medicine, the constituents isolated from its leaves were tested against MCF-7 human breast cancer cells. Of the compounds tested (Table 6), flemingins A (1) and C (3) exhibited cytotoxic activity. Although these compounds may be of interest for the development of novel anticancer agents, this observation indicates a possible risk of using the leaves of *F.*

Figure 2. Superimposed ^1H NMR spectra (DMSO- d_6 , 800.09 MHz, 25 $^\circ\text{C}$) of compounds **10** (red) and **11** (blue).

Figure 3. Key NOE correlations (blue arrows) observed for compound **12** (mixing time 700 ms, CD_3CN , 25 $^\circ\text{C}$).

Table 6. Cytotoxic (IC_{50} μM) and Radical Scavenging (ED_{50} μM) Activities of Selected Flemingins and of the Crude Leaf Extract of *Flemingia grahamiana*

sample	IC_{50}^b	ED_{50}^c	activity index
<i>F. grahamiana</i> (leaf extract) ^a		5.9	
flemingin A (1)	8.9	4.4	1.8
flemingin B (2)	>236.7	7.8	3.1
flemingin C (3)	7.6	5.9	2.4
flemingin G (4)	220.8	8.3	3.3
flemingin H (5)		8.9	3.5
flemingin I (6)	>236.3		
flemingin K (8)	>227.7		
flemingin L (9)	>236.3		
emodin (14)	317.4		
quercetin (standard)		2.5	1.0

^aThe chlorophyll was removed from the leaves by column chromatography on Sephadex LH-20 with CH_2Cl_2 – CH_3OH , 1:1, before testing. ^bAs positive control 1-isopropyl-3-(pyridin-4-ylethynyl)-1H-pyrazolo[3,4-*d*]pyrimidin-4-amine (IC_{50} = 5.0 μM , confidence interval (95%) = 1.4–17.8 nM)¹⁰ was used. ^c ED_{50} is given in $\mu\text{g}/\text{mL}$ for crude and in μM for pure compounds. The generally accepted threshold for bioactivity is 10 μM .

grahamiana by indigenous populations as a purgative and against skin diseases.²

The crude leaf methanol extract of *F. grahamiana* showed antioxidant activity in the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay.⁹ Strong antioxidant activities were observed for flemingins A–C (**1**–**3**), G (**4**), and H (**5**) (Table 6). The 2',5'-*para*-dihydroxylation of the B-ring is suggested to contribute to the strong antioxidant activity of these flemingins due to the formation of a stable 1,4-quinone upon oxidation. Despite the fact that its A-ring is not *para*-dihydroxylated, unlike **3** and **4**, providing the theoretical ability of radical scavenging by formation of a second quinoid ring, the most potent antioxidant activity among the tested compounds was observed for flemingin A (**1**).

EXPERIMENTAL SECTION

General Experimental Procedures. UV spectra were obtained on a Hewlett-Packard 8453 spectrophotometer; CD spectra were recorded on a JASCO J-710 CD spectropolarimeter. NMR spectra were acquired on Varian Unity Inova spectrometers operating at 200, 400, 600, and 800 MHz or on a Bruker 300 MHz spectrometer. NMR spectra were processed using MestReNova-9.0. Structural assignment was based on ^1H and ^{13}C NMR as well as gCOSY,¹¹ gNOESY,¹² gHSQC,¹³ and gHMBC¹⁴ spectra. The solvent residual peak was used for chemical shift referencing (DMSO- d_6 , δ_{H} 2.50 and δ_{C} 39.52; CD_3OD , δ_{H} 3.31 and δ_{C} 49.00; CD_3CN , δ_{H} 1.94 and δ_{C} 1.32). LC-MS (ESI) chromatograms were acquired on a PerkinElmer PE SCIEX API 150EX instrument equipped with a Turbolon spray ion source and a Gemini 5 mm RPC_{18} 110 Å column and applying a H_2O – CH_3CN , 80:20–20:80, gradient solvent system with a separation time of 8 min. HRESIMS were obtained with a Q-TOF-LC/MS spectrometer (Stenhagen Analyslab AB, Gothenburg, Sweden) using a 2.1 $\times$ 30 mm, 1.7 μm RPC_{18} column and a H_2O – CH_3CN gradient system (5:95–95:5 gradient and 0.2% formic acid). Column chromatography (CC) was carried out using silica gel 60 (70–230 mesh). Analytical thin-layer chromatography (TLC) was performed on silica gel 60 F_{254} (Merck) precoated aluminum plates, which after development with an appropriate solvent system were evaluated under UV light (254 and 366 nm). Gel filtration was carried out over Sephadex LH-20 (Pharmacia) suspended in CH_2Cl_2 – CH_3OH (1:1). Preparative HPLC was performed on a Waters 600E system using the Chromulan (Pikron Ltd.) software and an RP- C_8 Kromasil column (250 mm $\times$ 25 mm) with the solvent system H_2O – CH_3OH or H_2O – CH_3CN (gradient, 90:10 to 5:95, for 30–80 min, flow rate of 8–15 mL/min).

Plant Material. The leaves of *Flemingia grahamiana* were collected from Kitale in Western Kenya in October 2008. The plant material was identified by Mr. Simon Mathenge, a senior botanist at the East African Herbarium, Nairobi, where a voucher specimen (Mathenge 2008/487) is deposited.

Extraction and Isolation. Dried and pulverized leaves (413 g) of *F. grahamiana* were soaked twice in CH_2Cl_2 – CH_3OH (1:1) at room temperature, for 24 h in each case, to yield 30 g of crude extract after solvent evaporation. A portion of the extract (20 g) was adsorbed on silica gel 60 (70–230 mesh ASTM) (40 g), loaded onto a column of silica gel (38 cm length and 5 cm diameter, 200 g silica gel), and then eluted with increasing amounts of EtOAc in *n*-hexane to give 20 fractions, of approximately 500 mL each. Fractions 7–20 were purified by gel filtration over Sephadex LH-20 (eluted with CH_2Cl_2 – CH_3OH , 1:1). Fractions 7–9, eluted at 5–25% EtOAc in *n*-hexane, were combined and further purified by CC on silica gel deactivated with oxalic acid using an EtOAc in *n*-hexane gradient. This led to the isolation of compounds **1** (46.0 mg) and **14** (2.1 mg). Subfractions 7–9 were combined and subjected to preparative HPLC (RPC_8 250 mm $\times$ 25 mm, Kromasil column; H_2O – CH_3CN eluent), yielding compounds **12** (2.5 mg) and **13** (3.0 mg). Fraction 15 (eluted with 50% EtOAc in *n*-hexane) was subjected to preparative HPLC (H_2O – CH_3OH mixtures as eluents) and yielded compounds **4** (15.4 mg) and **5** (6.3 mg). Fractions 16 and 17, eluted with 50–70% EtOAc in *n*-hexane, were mixed and subjected to preparative HPLC (H_2O – CH_3OH as eluent) to give compound **8** (13.2 mg). Fractions 18 and 19, eluted with 70% to 100% EtOAc in *n*-hexane, were combined and separated on preparative HPLC (H_2O – CH_3OH eluent), giving a

mixed fraction (compounds **10** and **11**). Further purification of the latter by preparative HPLC with a gradient of H₂O–CH₃OH gave compounds **10** (1.5 mg) and **11** (2.3 mg). Fractions 13 and 12, eluted at 40% EtOAc in *n*-hexane, provided a crude mixture of **6**, **7**, and **9** by preparative HPLC (H₂O–CH₃CN eluent), which were then isolated using a gradient of H₂O–CH₃OH to give **3.0**, **2.6**, and **10.0** mg, respectively. The combined fractions **10**–**12**, eluted with 30% EtOAc in *n*-hexane, were subjected to repeated preparative HPLC (H₂O–CH₃CN eluent) chromatographic purification, yielding compounds **2** (20.6 mg) and **3** (30.7 mg).

Flemingin A (1): yellow solid; ¹H and ¹³C NMR data, see Tables S1 and S2 in the Supporting Information; ESIMS *m/z* 407.7 [M + H⁺]. The NMR and MS data were in agreement with the data given in ref 3.

Flemingin B (2): yellow solid; UV (MeOH) λ_{max} (log ε) 350 (4.3) nm; ESIMS *m/z* 423.1 [M + H⁺]. The ¹H and ¹³C NMR data, shown in the Supporting Information, were in agreement with the data given in ref 3.

Flemingin C (3): yellow solid; UV (MeOH) λ_{max} (log ε) 290 (4.2), 375 (4.1), 400 (4.2) nm; ¹H and ¹³C NMR data, see Supporting Information; ESIMS *m/z* 423.0 [M + H⁺]. The NMR and MS data were in agreement with the data given in ref 3.

Flemingin D (4): yellow, amorphous solid; UV (CH₃OH) λ_{max} (log ε) 295 (3.2) and 405 (3.1) nm; ¹H and ¹³C NMR data, see Tables 1 and 2; HRESIMS *m/z* 439.1771 [M + H]⁺ (calcd for C₂₅H₂₆O₇, 439.1757).

Flemingin H (5): yellow, amorphous solid; UV (CH₃OH) λ_{max} (log ε) 170 (3.3) and 365 (3.1) nm; ¹H and ¹³C NMR data, see Tables 1 and 2; HRESIMS *m/z* 439.1809 [M + H]⁺ (calcd for C₂₅H₂₆O₇, 439.1757).

Flemingin I (6): yellow, amorphous solid; UV (CH₃OH) λ_{max} (log ε) 275 (3.5), 365 (3.2) nm; CD no Cotton effect; ¹H and ¹³C NMR data, see Tables 1 and 2; HRESIMS *m/z* 423.1821 [M + H]⁺ (calcd for C₂₅H₂₆O₆, 423.1808).

Flemingin J (7): yellow, amorphous solid; (CH₃OH) λ_{max} (log ε) 305 (3.5), 340 (3.4), 370 (3.4) nm; CD no Cotton effect; ¹H and ¹³C NMR data, see Tables 1 and 2; HRESIMS *m/z* 423.1843 [M + H]⁺ (calcd for C₂₅H₂₆O₆, 423.1808).

Flemingin K (8): yellow, amorphous solid; ¹H and ¹³C NMR data, see Tables 3 and 4; HRESIMS *m/z* 439.1733 [M + H]⁺ (calcd for C₂₅H₂₆O₇, 439.1757).

Flemingin L (9): yellow, amorphous solid; UV (CH₃OH) λ_{max} (log ε) 265 (3.2), 285 (3.3), 300 (3.1) nm; ¹H and ¹³C NMR data, see Tables 3 and 4; HRESIMS *m/z* 423.1832 [M + H]⁺ (calcd for C₂₅H₂₆O₆, 423.1808).

Flemingin M (10): yellow, amorphous solid; UV (CH₃OH) λ_{max} (log ε) 265 (3.5), 380 (3.2) nm; CD no Cotton effect; ¹H and ¹³C NMR data, see Tables 3 and 4; HRESIMS *m/z* 441.1779 [M + H]⁺ (calcd for C₂₅H₂₈O₇, 441.1813).

Flemingin N (11): yellow, amorphous solid; UV (CH₃OH) λ_{max} (log ε) 275 (3.4), 350 (3.4) nm; CD no Cotton effect; ¹H and ¹³C NMR data, see Tables 3 and 5; HRESIMS *m/z* 441.1912 [M + H]⁺ (calcd for C₂₅H₂₈O₇, 441.1813).

Flemingin O (12): yellow, amorphous solid; UV (CH₃OH) λ_{max} (log ε) 300 (3.3), 360 (3.1) nm; CD no Cotton effect; ¹H and ¹³C NMR data, see Tables 3 and 5; HRESIMS *m/z* 407.1878 [M + H]⁺ (calcd for C₂₅H₂₇O₅, 407.1858).

Deoxyhomoflemingin (13): yellow, amorphous solid; UV (MeOH) λ_{max} (log ε) 300 (3.2) nm; ¹H and ¹³C NMR data see Supporting Information; ESIMS *m/z* 423.4 [M + H]⁺. The NMR data were in agreement with the data given in ref 4.

Cytotoxicity Assays. Cytotoxicity assays against MCF-7 human breast cancer cells were carried out as described by Endale et al.¹⁵ The cells were cultured in Dulbecco's modified Eagle's medium supplemented with fetal bovine serum (10% v/v), 2 mM L-glutamine, 100 units/mL penicillin, and 100 μg/mL streptomycin. The medium was preincubated at 37 °C under humid conditions with a mixture of 5% CO₂ and 95% air. Cells were seeded in 96-well plates at optimal cell density to ensure exponential growth for the duration of the assay. The growth medium was replaced with the experimental medium containing an appropriate drug concentration or control (0.1% or 1.0%

v/v, dimethyl sulfoxide) after 24 h of preincubation. Cell viability was measured, after 48 h of incubation, using PrestoBlue cell viability reagent (Invitrogen AB Lidingö, Sweden), in accordance with the manufacturer's instructions. The fluorescence was measured using a POLARstar Omega (BMG Labtech, Ortenberg, Germany) plate reader. Results are expressed as means ± SE for six replicates as a percentage of the vehicle control (assumed to be 100%). Experiments were carried out independently at least three times. Statistical analyses were performed using a two-tailed Student's *t*-test. A value of *p* < 0.05 was considered to indicate statistically significant differences.

Antioxidant Activity Assays. The method described by Ohnishi et al.¹⁶ was used with minor modifications. Test sample solutions at concentrations of 320, 160, 80, 40, 20, 10, 5, and 2.5 μg/mL were prepared in double-distilled methanol. From each concentration, 1.0 mL of the sample was added to 2.0 mL of 76 μM DPPH dissolved in methanol. The mixture was allowed to stand at room temperature for 30 min, and the absorbance of the remaining DPPH was then measured at 517 nm. The radical scavenging activity was measured as the decrease in absorbance due to the DPPH radical, expressed as a percentage of the absorbance of the control solution (1.0 mL of methanol and 2.0 mL of 76 μM DPPH solution). Activity was then expressed as EC₅₀, the concentration of the test compound required to give a 50% decrease in the absorbance compared to that of the control solution. Quercetin was used as the positive control. The percentage of scavenged DPPH was calculated as 100 × (A_{DPPH} – A_{Sample})/A_{DPPH}, where A_{DPPH} is the absorbance of the solution containing DPPH but without a test sample and A_{Sample} is the absorbance of the mixture of the test sample and DPPH solution. Values of EC₅₀ were determined by plotting the percentage of scavenged DPPH versus the initial concentration of each sample. The activity index was calculated by comparing the value of EC₅₀ (in μM) for a given test compound with that of the standard.

■ ASSOCIATED CONTENT

● Supporting Information

1D and 2D NMR, MS, UV, and CD spectra and data for cytotoxicity and antioxidant assays. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Authors

*(A. Yenesew) Tel: +254 733 832 576. Fax: +254 20 444 6138. E-mail: ayenesew@uonbi.ac.ke.

*(M. Erdélyi) Tel: +46 31 786 9033. E-mail: mate@chem.gu.se.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful to S. Mathenge of the East African Herbarium, Nairobi, Kenya, for identifying the plant species studied. I.G. is grateful to the Swedish Institute for a research fellowship and to the German Academic Exchange Services (DAAD) for a Ph.D. scholarship, which was offered through the Natural Products Research Network for Eastern and Central Africa (NAPRECA). The Swedish Institute, the Swedish Research Council (2012-6124), and Sigrud and Elsa Golje's Memorial Foundation (2013-0298) are acknowledged for funding this research.

■ REFERENCES

- (1) Gillet, J. B.; Polhill, B. A.; Verdcourt, B. Leguminosae. In *Flora of Tropical East Africa*; Milne-Redhead, E., Polhill, R. M., Eds.; Crown Agents for Overseas Governments and Administrations: London, 1971; pp 1–501.
- (2) Gahlot, K.; Lal, V. K.; Jha, S. *Int. J. Phytomed.* **2011**, *3*, 294–307.

- (3) Cardillo, G.; Merlini, L.; Mondelli, R. *Tetrahedron* **1968**, *24*, 497–510.
- (4) Cardillo, B.; Gennaro, A.; Merlini, L.; Nasini, G.; Servi, S. *Phytochemistry* **1973**, *12*, 2027–2031.
- (5) Cardillo, G.; Merlini, L.; Mondelli, R.; Selva, A. *Ric. Sci.* **1967**, *37*, 826–828.
- (6) Danielsen, K.; Aksnes, D. W. *Magn. Reson. Chem.* **1992**, *30*, 359–360.
- (7) Prokofev, E. P.; Grigoreva, N. Y.; Semenovskii, A. V. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **1980**, *29*, 586–589.
- (8) Dewick, P. M. *Medicinal Natural Products: A Biosynthetic Approach*, 2nd ed.; John Wiley & Sons: Chichester, UK, 2002; pp 140–151.
- (9) Sharma, O. P.; Bhat, T. K. *Food Chem.* **2009**, *113*, 1202–1205.
- (10) Diner, P.; Alao, J. P.; Söderlund, J.; Sunnerhagen, P.; Grötl, M. *J. Med. Chem.* **2012**, *55*, 4872–4876.
- (11) Wokaun, A.; Ernst, R. R. *Chem. Phys. Lett.* **1977**, *52*, 407–412.
- (12) Kumar, A.; Ernst, R. R.; Wuthrich, K. *Biochem. Biophys. Res. Commun.* **1980**, *95*, 1–6.
- (13) Perpichdumont, M.; Reynolds, W. F.; Enriquez, R. G. *Magn. Reson. Chem.* **1988**, *26*, 881–887.
- (14) Parella, T. *Magn. Reson. Chem.* **1998**, *36*, 467–495.
- (15) Endale, M.; Ekberg, A.; Akala, H. M.; Alao, J. P.; Sunnerhagen, P.; Yenesew, A.; Erdelyi, M. *J. Nat. Prod.* **2012**, *75*, 1299–1304.
- (16) Ohnishi, S. T.; Kojima, R. *FASEB J.* **1994**, *8*, A596–A596.