



## CASE REPORT

### Management of Puff Adder (*Bitis Arietans*) Snake Bite and Envenomation in Dogs: Case Report

WE Mwangi\*, P Kimeli, LW Mathai, DW Muasya and AN Kipyegon

Department of Clinical Studies, Faculty of Veterinary Medicine, University of Nairobi P.O Box 29053 – 00625, Kangemi, Kenya

#### ARTICLE INFO

Received: July 01, 2014

Revised: July 09, 2014

Accepted: July 13, 2014

#### Key words:

Antihistamines

Anti-snake venom

Dog

Snake bite

#### \*Corresponding Author

WE Mwangi

drwillymwangi@gmail.com

#### ABSTRACT

Two mongrel dogs were presented with a history of recumbency, salivation, dullness and depression that had started following snake bites. Physical examination revealed swellings with fang marks and hemorrhage in the area of bites. The hematological parameters showed reduced values of erythrocyte counts, platelet counts, hemoglobin, packed cell volume and increased total leukocyte count. The biochemical values showed elevated levels of alanine aminotransferase and creatinine. The successful treatment was done with anti-snake venom, fluids, corticosteroid, antihistamines and antibiotic.

Cite This Article as: Mwangi WE, P Kimeli, LW Mathai, DW Muasya and AN Kipyegon, 2014. Management of puff adder (*Bitis arietans*) snake bite and envenomation in dogs: case report. Inter J Vet Sci, 3(3): 160-162.

www.ijvets.com